Florida Citrus Advanced Technology Program

QUARTERLY & FINAL REPORTS: Control of Citrus Greening, Canker & Emerging Diseases of Citrus

Instructions Complete the fields based on your project specs. When finished, save the form to your local disk using a unique name. Then, go to **http://research.fcprac.com**, and log in with your user name and password using Researcher Login in the lower left. Find this project title and click on **Submit a Report**. Update your profile information if needed, then upload this report as directed.

2009-2010 REPORT	CATEGORY (drop-down)	TODAY'S DATE (m/d/yr)
○ *Quarterly Report* ◉ *Annual Report* ○ *Final*	Psyllid	6/15/10

WHAT IS THE "HEADLINE" FOR THIS REPORT (e.g. a one-sentence "newspaper headline" describing what you accomplished)

Systemic insecticides and mineral oil interferes with the ACP feeding behavior and can prevent the bacteria transmission.

TITLE and CONTACT INFORMATION

Proposal Title CAN INSECTICIDES AND MINERAL OIL AVOID TRANSMISSION OF Candidatus Liberibacter asiaticus BY Diaphorina citri?

Principal Investigator Pedro Takao Yamamoto	*PI Last Name* Yamamoto
Email ptyamamoto@fundecitrus.com.br	*FDACS Contract Number* NAS 236
Phone +55 16 33017025	*Project Duration (years)* 2 *Year of Project* 2
Organization Fundecitrus	*Total Direct Funds (current year)* US$ 17,710.00

REPORT UPDATE (650 words; provide details about your headline)

Imidacloprid (Confidor 700 GrDA), 0.35 g AI/plant and thiamethoxam (Actara 250 WG) 0.25 g AI/plant, applied in the nursery tree bags, before planting, was efficient to control ACP until 60 days after application. The time to cause 100% of ACP mortality was between 5 to 7 days after the confinement of adults in treated plants. However, researches using electrical penetration graph (EPG) showed that in plants treated with imidacloprid and thiamethoxam, after the first feeding on phloem, the adults do not do more probing. We carried out the first PCR of the plants in this experiment and the results were negative, no plants have been detected the presence of the bacterium L. Ca asiaticus. No transmission results yet. We finish the second experiment that was performed to determine if the systemic insecticides are effective until 90 days after application and its effect on transmission of the bacteria. In this experiment, the time to reach 100% of mortality ranged from 3 to 7 days for both systemic insecticides tested (imidacloprid and thiamethoxam). The insecticides were effective up to 90 days after application. The results of PCR carried out for the ACP, in some periods, were positive for 100% of the samples, consisting of 10 insects tested, but in the confinement held at 46 days after application, in any sample was detected the presence of the bacteria. No acquisition in this period. In bioassays performed at 75 and 90 days after application, the percentage of positive samples was 50 to 70% and 10 to 40%, respectively.

We started the experiment 2, the difference from the experiment 1 is the application of varying doses of the systemic insecticides and confinement of the ACP in plants treated only 7 days after application. To thiamethoxam (Actara 250 WG), the doses tested were: 1, 0.5, 0.1 and 0.05 g/nursery tree and imidacloprid (Provado 200 SC) were: 1.75, 0.9, 0.2 and 0.08 mL/nursery tree. We also started the experiment 3, using different insecticide spraying to determine if they prevent the transmission and for how long.

Plants treated with insecticide, the proportion of insects reaching the phloem was similar between plants treated with imidacloprid (0.35 g AI/tree), thiamethoxam (0.25 g AI/tree) and control (untreated plants), being respectively 74, 72 and 76%. The time to perform the first ACP salivation was also similar between treatments, 118.4, 103.2, and 112.6 minutes, respectively. However, the time of phloem ingestion is drastically reduced compared to untreated plants: imidacloprid 6.1, 9.9 and 6.9 min, respectively for 15, 35 and 95 days after application (DAA); thiamethoxam 9.6, 14.5 and 17.5 min, respectively for 15, 35 and 95 DAA; Control 142.0, 80.3 and 129.0 minutes, respectively for 15, 35 and 95 DAA. Apparently, ACP can only distinguish between plants with and without treatment from the moment that start ingesting the phloem sap. In this case, it was observed that after ingestion of sap with insecticide, the ACP removes the stylet from the plant and rarely returns to start a new probe on the same plant.

In plants sprayed with mineral oil decreased the percentage of psyllids that can reach the phloem when compared with plants not sprayed, 20 and 70% respectively. However, the few insects that reach the phloem of treated plants carry out long periods of ingestion in this vascular tissue (≥ 1 h).

We began the experiment to evaluate the effect of oil on the feeding behavior of ACP and its effect on repellency of the vector. The results showed that up to 21 days after application, mineral oil, 1.5%, shows repellency to adults of D. citri. Using electrical penetration graphs (EPG) techniques, we are studying the probing behavior of ACP in plants that were applied mineral oil. The results showed that the number of insects reaching the phloem is lower. The next step is the study of lower doses of mineral oil.